


Malaria




**2014 Kenya Demographic
and Health Survey (KDHS)**

*Follow along on
Twitter!
#KenyaDHS
@KNBS2013
@DHSprogram*


Ownership of Mosquito Nets

Percent of households with at least one mosquito net




Use of ITNs

Percent who slept under an ITN the night before the survey among all households




Percent who slept under an ITN the night before the survey among households with at least one ITN



Trends in Use of ITNs


Percent of children under 5 and pregnant women age 15-49 who slept under an ITN the night before the survey

■ 2003 KDHS ■ 2008-09 KDHS ■ 2014 KDHS



Intermittent Preventive Treatment during Pregnancy (IPTp)

Percent of women age 15-49 with a live birth in the two years before the survey who:




Fever in Children

In the 2 weeks before the survey, **24%** of children under 5 had a **fever**.


- **72%** were taken to a health facility or provider for **advice or treatment**.
- **35%** had a **blood taken from finger or heel for testing**.
- **27%** were given **antimalarial drugs**.

Type of Antimalarial Drug by Children

Among children under 5 with fever in the two weeks before the survey who took an antimalarial, percent who took a type of antimalarial drug



HIV Knowledge, Attitudes, and Behaviour




**2014 Kenya Demographic
and Health Survey (KDHS)**

*Follow along on
Twitter!
#KenyaDHS
@KNBS2013
@DHSprogram*

Knowledge of HIV Prevention Methods by Education

Percent of women and men age 15-49 who know that HIV can be prevented by using condoms AND limiting sex to one uninfected partner

■ Women ■ Men



Knowledge of HIV Prevention Methods by County

**< 50% know that HIV can be prevented
by using condoms AND limiting sex to one uninfected
partner**

- Kilifi Men (44%)
- Garissa Women (31%)
- Wajir Women (15%); Men (11%)
- Mandera Women (13%)
- Turkana Women (49%); Men (2%)
- West Pokot Women (48%)


Multiple Sexual Partners

■ Women ■ Men

Among women and men age 15-49 who had 2+ sexual partners in past 12 months, percent who:

Among women and men age 15-49 who have ever had sexual intercourse:

Percent of women and men age 15-49 who had:




91% of women and **97%** of men
know where to get an HIV test.

HIV Testing

Percent of women and men age 15-49

■ Women ■ Men




HIV Testing during Pregnancy

Among women who gave birth in the two years before the survey, **67%** were **counselled and tested for HIV** during antenatal care and **received the results.**

Trends in Male Circumcision


Percent of men age 15-49 who report having been circumcised



Comprehensive Knowledge of HIV among Youth by Residence

Percent of women and men age 15-24 with comprehensive knowledge of HIV*

■ Women ■ Men




*Comprehensive knowledge means knowing that the risk of getting HIV can be reduced by using condoms and limiting sex to one uninfected partner, knowing that a healthy looking person can have HIV, and rejecting the two most common local misconceptions about HIV prevention and transmission.

Knowledge of a Condom Source among Youth by Residence

Percent of women and men age 15-24 who know a condom source

■ Women ■ Men



Age at First Sexual Intercourse


- **12%** of young women and **21%** of young men age 15-24 had sexual intercourse **before age 15.**
- **47%** of young women and **55%** of young men age 18-24 had sexual intercourse **before age 18.**

Premarital Sexual Intercourse and Condom Use among Youth

■ Women ■ Men

Percent of never-married women and men age 15-24 who:

Among never-married women and men age 15-24 who had sexual intercourse in past 12 months, percent who:



HIV Testing among Youth

Among women and men age 15-24 who had sexual intercourse in past 12 months, percent tested for HIV in past 12 months and received the results

